

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner  
US Department of Commerce  
United States Patent and Trademark  
Office, PCT  
2011 South Clark Place Room  
CP2/5C24  
Arlington, VA 22202  
ETATS-UNIS D'AMERIQUE  
in its capacity as elected Office

Date of mailing (day/month/year) 23 April 2001 (23.04.01)	
International application No. PCT/US00/20641	Applicant's or agent's file reference 15280-4003PC
International filing date (day/month/year) 27 July 2000 (27.07.00)	Priority date (day/month/year) 28 July 1999 (28.07.99)
Applicant FRANCHINI, Genoveffa et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

12 February 2001 (12.02.01)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer Juan Cruz
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38

## PATENT COOPERATION TREATY

## PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

REC'D 05 NOV 2001

WIPO PCT

Applicant's or agent's file reference 15280-4003PC	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US00/20641	International filing date (day/month/year) 27/07/2000	Priority date (day/month/year) 28/07/1999
International Patent Classification (IPC) or national classification and IPC A61K39/00		
Applicant THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 5 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 12/02/2001	Date of completion of this report 02.11.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office - P.B. 5818 Patentlaan 2 NL-2280 HV Rijswijk - Pays Bas Tel. +31 70 340 - 2040 Tx: 31 651 epo nl Fax: +31 70 340 - 3016	Authorized officer Skelly, J Telephone No. +31 70 340 3413 

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US00/20641

## I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

**Description, pages:**

1-26 as originally filed

**Claims, No.:**

1-17 as originally filed

**Drawings, sheets:**

1/8-8/8 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
  - ☐ the language of publication of the international application (under Rule 48.3(b)).
  - ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).
3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:
- ☐ contained in the international application in written form.
  - ☐ filed together with the international application in computer readable form.
  - ☐ furnished subsequently to this Authority in written form.
  - ☐ furnished subsequently to this Authority in computer readable form.
  - ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
  - ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US00/20641

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes: Claims
	No: Claims 1-17
Inventive step (IS)	Yes: Claims
	No: Claims 1-17
Industrial applicability (IA)	Yes: Claims
	No: Claims 1-17 see separate sheet

2. Citations and explanations  
**see separate sheet**

## VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:  
**see separate sheet**

**R Item V**

**Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

Reference is made to the following documents:

D1: WO 98 08539 A (CHIRON CORPORATION) 5 March 1998 (1998-03-05)

D2: Z. HEL ET AL.: 'Viremia control following antiretroviral treatment and therapeutic immunisation during primary SIV251 infection of macaques', NATURE MEDICINE, vol. 6, 2000, pages 1140-1146

**Novelty (Art. 33(2) PCT)**

D1 discloses a therapeutic regimen for HIV-infected individuals comprising reducing viral load by administration of a viral load reducer, such as a combination of reverse transcriptase and protease inhibitors (page 21, lines 7-10; page 45, line 10- page 46, line 9), increasing the level of CD4+ cells, for example with IL-2 (page 46, lines 10-27) and administering a vaccine to potentiate a cytotoxic of CD8+ response, which may be nucleic acid based (page 50, line 9-page 51, line 18), and expressed from poxviral vectors (page 6, lines 4-16). These steps may be simultaneous or sequential (page 69, lines 2-10). The subject-matter of claims 1-17 is therefore not novel.

**Inventive Step (Art. 33(3) PCT)**

The problem stated to be solved by the invention is to provide a means of stimulating a protective CD8+-mediated immune response against HIV in infected persons. Before an inventive step can be recognised it must be clear that this problem has been solved by the claimed method. The publication for the application (D2) shows that vaccination made no difference in viral rebound after cessation of treatment despite increasing CD4+ and CD8+ responses (see page 1142, col. 2, -page 1143, col.1, line 3) Hence a "protective" immune response as claimed has not been achieved and the subject-matter of claims 1-17 lacks inventive step.

**Industrial Applicability (Art. 33(4) PCT)**

For the assessment of the present claims 1-17 on the question whether they are industrially applicable, no unified criteria exist in the PCT Contracting States. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject-matter of claims to the use of a compound in medical treatment, but may allow, however, claims to a known compound for first use in medical treatment and the use of such a compound for the manufacture of a medicament for a new medical treatment.

**Re Item VIII**

**Certain observations on the international application**

The invention is defined by a result to be achieved: "A method of stimulating an efficient CD8+

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US00/20641

response...where said peptides are presented **in an amount sufficient to stimulate a protective CD8+ response...**". This broad formulation merely amounts to claiming the underlying technical problem. There is no evidence in the description that the applicant has solved this problem, on the contrary (see D2, loc. cit.). Hence the description does not disclose the invention in a manner sufficiently clear and complete for it to be carried out by a person skilled in the art (Art. 5 PCT). The subject-matter of the claims is therefore not fully supported by the description (Art. 6 PCT).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number  
**WO 01/08702 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 39/00**

20817 (US). TARTAGLIA, James [US/US]; 7 Christina Drive East, Schenectady, NY 12303 (US). NACSA, Janos [HU/US]; 10834 Margate Road, Silver Spring, MD 20901 (US).

(21) International Application Number: **PCT/US00/20641**

(22) International Filing Date: **27 July 2000 (27.07.2000)**

(25) Filing Language: **English**

(74) Agents: **LOCKYER, Jean, M. et al.**; Townsend and Townsend and Crew LLP, 8th Floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).

(26) Publication Language: **English**

(30) Priority Data:

60/146,240	28 July 1999 (28.07.1999)	US
60/178,989	28 January 2000 (28.01.2000)	US
60/200,445	28 April 2000 (28.04.2000)	US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **THE GOVERNMENT OF THE UNITED STATES OF AMERICA**, represented by **THE SECRETARY OF THE DEPARTMENT OF HEALTH AND HUMAN SERVICES** [US/US]; Bethesda, MD 20892 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRANCHINI, Genoveffa** [US/US]; 4400 17th Street N.W., Washington, DC 20011 (US). **HEL, Zdenek** [CZ/US]; 11120 Troy Road, North Bethesda, MD 20852 (US). **SHEARER, Gene** [US/US]; 5512 Glenwood Road, Bethesda, MD


Published:

— Without international search report and to be republished upon receipt of that report.

[Continued on next page]

(54) Title: **IMMUNOTHERAPY IN HIV INFECTED PERSONS USING VACCINES AFTER MULTI-DRUG TREATMENT**

**VIRAL LOAD AND CD4<sup>+</sup> T-CELL RESPONSES  
IN UNTREATED/TREATED NYVAC-SIV VACCINATED MACAQUES**



(57) Abstract: This invention relates to an improved method of maintaining an immuno-protective response in persons infected with a retrovirus after highly active anti-retroviral therapy.

WO 01/08702 A2



*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

# **IMMUNOTHERAPY IN HIV INFECTED PERSONS USING VACCINES AFTER MULTI-DRUG TREATMENT**

5

## **CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims priority to U.S. Provisional Application Serial Nos. 60/146,240, filed July 28, 1999; 60/178,989, filed January 28, 2000; and 60/200,445, filed  
10 April 28, 2000, each of which is incorporated by reference herein.

## **STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT**

[ Not Applicable ]

15

## **FIELD OF THE INVENTION**

This invention relates to an improved method of maintaining an immuno-  
protective response in persons infected with a retrovirus after highly active anti-  
20 retroviral therapy (HAART). Surprisingly, these patients demonstrate a CD8<sup>+</sup>  
response following HAART treatment.

## **SUMMARY OF THE INVENTION**

25 The present invention is directed to a method of stimulating an efficient CD8<sup>+</sup>  
response in a human infected with an HIV or HTLV-1 retrovirus, who has a viral load  
of less than 10,000 viral copies, often 5,000 or 2,000 or less, per ml of plasma and a  
CD4<sup>+</sup> cell count that is often above 500 cells/ml, but can be above 400 cells/ml or 300  
cells/ml; and who has been treated with one or more anti-viral agents, which  
30 contributed to a lower viral copy and higher CD4<sup>+</sup> cell count than before treatment.  
The method comprises administering a nucleic acid-based vaccine, which enters the  
cells and intracellularly produces HIV- or HTLV-1-specific peptides for presentation  
on the cell's MHC class I molecules in an amount sufficient to stimulate a protective  
CD8<sup>+</sup> response.

In a preferred embodiment the human has been treated with anti-viral agents, which resulted in the human having a viral load of less than 1,000 viral copies per ml of blood serum and a CD4<sup>+</sup> cell count that is above 500 cells/ml. The anti-viral agents can preferably comprise a combination of inhibitors of proteases and inhibitors of reverse transcriptase.

The method can use a vaccine that is a DNA based vaccine or that is an attenuated recombinant virus. A preferred virus is an attenuated pox virus, particularly NYVAC and ALVAC, attenuated vaccinia and canarypox viruses respectively. Other attenuated pox viruses such as MVA can also be used.

The vaccine can further comprise an adjuvant and may be administered a second time. The vaccine can also comprise interleukin-2 (IL-2) and/or CD40 ligand in an amount that is sufficient to potentiate the CD8<sup>+</sup> response.

The method of the invention can be particularly useful for a person who has been infected with HIV and has demonstrated repeated and sustained proliferative T-cell responses to gp120 envelope protein or both gp 120 envelope and p24 gag antigen.

The person infected with HIV can be further tested by a skin test for a hypersensitive response to the p24 gag antigen or to the gp120 envelope antigen.

## BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Figure 1 shows the viral load and proliferative CD4<sup>+</sup> T-helper responses to p27 *gag* and gp120 in the infected, treated vaccinated macaques. The top panels display the kinetics of virus load in the plasma animals from groups A, B, and C. In group A two animals (641 and 642) did not respond to therapy and were not included in the analyses.

Figure 2: Figure 2 shows the CD8<sup>+</sup> response in the infected, treated, vaccinated animals.

Figure 3: Figure 3 shows p27-specific (Fig. 3a and 3c) and gp120-specific (Fig. 3b and 3d) T-cell proliferation in SIV-infected HAART-treated *Rhesus macaques* following administration of a single dose of ALVAC-SIV<sub>gpe</sub>. (Fig. 3a and Fig. 3b) or

ALVAC (Fig. 3c and 3d). The arrows indicate time after SIV infection at which the animals were inoculated with vaccine.

Figure 4: Figure 4 shows the CD3<sup>+</sup>CD8<sup>+</sup> T-cell responses detected in fresh peripheral blood mononuclear cells from SIV-infected, HAART-treated macaques inoculated with ALVAC-SIV<sub>gpc</sub> (Fig. 4a) or ALVAC (Fig. 4b).

Figure 5: Figure 5 shows the induction of CD8<sup>+</sup> (Fig. 5a and 5b) and CD4<sup>+</sup> (Fig. 5c and 5d) T-cell responses following administration of a DNA vaccine to naïve macaques. DNA was administered at the times indicated by the arrow. The plasmids employed were pCMV-*gag* and pCMV-*env*, which are CMV expression plasmids expressing the *gag* and *env* genes, respectively.

Figure 6: Figure 6 shows the induction of CD8<sup>+</sup> (Fig. 6a and 6b) and CD4<sup>+</sup> (Fig. 6c and 6d) T-cell responses following administration of two inoculation of NYVAC-SIV-*gag-pol-env* to naïve macaques. The vaccine was at the times indicated by the arrows.

Figure 7: Figure 7 shows the administration schedule of NYVAC vaccine with and without interleukin-2 (IL-2) in HAART-treated macaques.

Figure 8: Figure 8 shows the amount of viral RNA (top panel) and the CD8<sup>+</sup> and CD4<sup>+</sup> proliferative responses, top and lower panel respectively, in a NYVAC-SIV-inoculated Rhesus macaque treated with IL-2. The horizontal bars indicate the length of treatment with anti-retroviral therapy (ART) and IL-2. Viral RNA levels in the plasmid (top panel) are indicated with open circles. The percent of CD8<sup>+</sup>CD3<sup>+</sup> tetramer-binding cells is indicated in the top panel by the hatched vertical bars. ELISPOT assay results measuring the release of  $\gamma$ -interferon are shown by the black vertical bars. Proliferative responses to p27 *gag* and gp120 are indicated in the lower panel. IL-2 was administered daily by subcutaneous injection at a dose of 120,000 units.

**Definitions**

“Attenuated recombinant virus” refers to a virus that has been genetically altered by modern molecular biological methods, *e.g.* restriction endonuclease and ligase treatment, and rendered less virulent than wild type, typically by deletion of specific genes or by serial passage in a non-natural host cell line or at cold temperatures.

“Efficient CD8<sup>+</sup> response” is referred to as the ability of cytotoxic CD8<sup>+</sup> T-cells to recognize and kill cells expressing foreign peptides in the context of a major histocompatibility complex (MHC) class I molecule.

“Nonstructural viral proteins” are those proteins that are needed for viral production but are not necessarily found as components of the viral particle. They include DNA binding proteins and enzymes that are encoded by viral genes but which are not present in the virions. Proteins are meant to include both the intact proteins and fragments of the proteins or peptides which are recognized by the immune cell as epitopes of the native protein.

“Nucleic acid-based vaccines” include both naked DNA and vectored DNA (within a viral capsid) where the nucleic acid encodes B-cell and T-cell epitopes and provides an immunoprotective response in the person being vaccinated.

“Plasma” refers to the fraction of whole blood resulting from low speed centrifugation of EDTA- or heparin- treated blood.

“Pox viruses” are large, enveloped viruses with double-stranded DNA that is covalently closed at the ends. Pox viruses replicate entirely in the cytoplasm, establishing discrete centers of viral synthesis. Their use as vaccines has been known since the early 1980’s (*see, e.g.* Panicali, D. *et al.* “Construction of live vaccines by using genetically engineered pox viruses: biological activity of recombinant vaccinia virus expressing influenza virus hemagglutinin”, *Proc. Natl. Acad. Sci. USA* 80:5364-5368, 1983).

A “retrovirus” is a virus containing an RNA genome and an enzyme, reverse transcriptase, which is an RNA-dependent DNA polymerase that uses an RNA molecule as a template for the synthesis of a complementary DNA strand. The DNA form of a retrovirus commonly integrates into the host-cell chromosomes and remains part of the host cell genome for the rest of the cell’s life.

“Structural viral proteins” are those proteins that are physically present in the virus. They include the capsid proteins and enzymes that are loaded into the capsid

with the genetic material. Because these proteins are exposed to the immune system in high concentrations, they are considered to be the proteins most likely to provide an antigenic and immunogenic response. Proteins are meant to include both the intact proteins and fragments of the proteins or peptides which are recognized by the immune cell as epitopes of the native protein.

“Viral load” is the amount of virus present in the blood of a patient. Viral load is also referred to as viral titer or viremia. Viral load can be measured in variety of standard ways.

## DETAILED DESCRIPTION

### Introduction

This invention is a novel therapeutic modality for treating persons infected with a lymphotropic or immune destroying retrovirus. A physician presented with a patient whose immune system is compromised by retroviral infection can elect to treat that patient with a host of powerful antiviral agents including inhibitors of viral proteases and reverse transcriptase. This is known as highly active anti-retroviral therapy (HAART). The conventional HAART protocols are complex and difficult for patients to follow. The drugs also have a number of problematic side effects. In addition, these expensive and complicated treatments do not eliminate the virus, but merely hold the virus in check. If the patient is non-compliant, the viral counts rebound. Accordingly, for the vast majority of patients, a lifetime of drugs is advised.

This invention is the discovery that after HIV infection, HAART treatment can sufficiently restore a patient’s immune system to effectively mount a CD8<sup>+</sup> response when a patient is provided with a CD8<sup>+</sup>-inducing vaccine. This response can effectively maintain a low titer of virus and significantly reduce the patient dependency on HAART when the CD8<sup>+</sup>-inducing vaccine is an HIV vaccine.. While some such vaccines have been suggested as useful for seropositive patients (U.S. Patent No. 5,863,542 column 18, lines 60-63), they are surprisingly effective for this subpopulation of seropositive patients and not for other seropositive patients.

### Vaccines of use in this invention

Vaccines useful for the induction of CD8<sup>+</sup> T-cell responses comprise nucleic acid-based vaccines (delivered via a viral vector or directly as a DNA vaccine) that

provide for the intracellular production of viral-specific peptide epitopes that are presented on MHC Class I molecules and subsequently induce an immunoprotective cytotoxic T lymphocyte (CTL) response.

The invention contemplates single or multiple administrations of a nucleic acid-based vaccine as a direct DNA vaccine or as a recombinant virus vaccine, or both. This vaccination regimen may be complemented with administration of recombinant protein vaccines (*infra*), or may be used with additional vaccine vehicles.

#### *Attenuated recombinant viral vaccines*

Attenuated recombinant viruses that express retrovirus specific epitopes are of use in this invention. Attenuated viruses are modified from their wildtype virulent form to be either symptomless or weakened when infecting humans. Among the recombinant viruses of use are adenoviruses, adeno-associated viruses, retroviruses and poxviruses.

A recombinant, attenuated virus for use in this invention as a vaccine is a virus wherein the genome of the virus is defective with respect to a gene essential for the efficient production of, or essential for the production of, infectious virus. The mutant virus acts as a vector for an immunogenic retroviral protein by virtue of the virus encoding foreign DNA. This provokes or stimulates a cell-mediated CD8<sup>+</sup> response.

The virus is then introduced into a human vaccinee by standard methods for vaccination of live vaccines. A live vaccine of the invention can be administered at, for example, about 10<sup>4</sup> -10<sup>8</sup> organisms/dose, or 10<sup>6</sup> to 10<sup>9</sup> pfu per dose. Actual dosages of such a vaccine can be readily determined by one of ordinary skill in the field of vaccine technology.

The selection of the virus is not critical. Examples of viral expression vectors include adenoviruses as described in M. Eloit *et al.*, "Construction of a Defective Adenovirus Vector Expressing the Pseudorabies Virus Glycoprotein gp50 and its Use as a Live Vaccine", *J. Gen. Virol.*, 71(10):2425-2431 (Oct., 1990).), adeno-associated viruses (*see, e.g.*, Samulski *et al.*, *J. Virol.* 61:3096-3101 (1987); Samulski *et al.*, *J. Virol.* 63:3822-3828 (1989)), papillomavirus, Epstein Barr virus (EBV) and Rhinoviruses (*see, e.g.*, U.S. Patent No. 5,714,374). Human parainfluenza viruses are also reported to be useful, especially JS CP45 HPIV-3 strain. The viral vector may be derived from herpes simplex virus (HSV) in which, for example, the gene encoding glycoprotein H (gH) has been inactivated or deleted. Other suitable viral vectors include retroviruses (*see, e.g.*, Miller, *Human Gene Ther.* 1:5-14 (1990); Ausubel *et al.*, *Current Protocols in Molecular Biology*).

The poxviruses are of preferred use in this invention. There are a variety of attenuated poxviruses that are available for use as a vaccine against HIV. These include attenuated vaccinia virus, cowpox virus and canarypox virus. In brief, the basic technique of inserting foreign genes into live infectious poxvirus involves a recombination between pox DNA sequences flanking a foreign genetic element in a donor plasmid and homologous sequences present in the rescuing poxvirus as described in Piccini *et al.*, *Methods in Enzymology* 153, 545-563 (1987). More specifically, the recombinant poxviruses are constructed in two steps known in the art and analogous to the methods for creating synthetic recombinants of poxviruses such as the vaccinia virus and avipox virus described in U.S. Pat. Nos. 4,769,330, 4,722,848, 4,603,112, 5,110,587, and 5,174,993, the disclosures of which are incorporated herein by reference.

First, the DNA gene sequence encoding an antigenic sequence, such as a known T-cell epitope, is selected to be inserted into the virus. The sequence is placed into an *E. coli* plasmid construct into which DNA homologous to a section of DNA of the poxvirus has been inserted. Separately, the DNA gene sequence to be inserted is ligated to a promoter. The promoter-gene linkage is positioned in the plasmid construct so that the promoter-gene linkage is flanked on both ends by DNA homologous to a DNA sequence flanking a region of pox DNA containing a nonessential locus. The resulting plasmid construct is then amplified by growth within *E. coli* bacteria.

Second, the isolated plasmid containing the DNA gene sequence to be inserted is transfected into a cell culture, *e.g.* chick embryo fibroblasts, along with the poxvirus. Recombination between homologous pox DNA in the plasmid and the viral genome respectively, gives a poxvirus modified by the presence, in a nonessential region of its genome, of foreign DNA sequences.

Attenuated recombinant pox viruses are a preferred vaccine. A detailed review of this technology is found in US Patent No. 5,863,542 which is incorporated by reference herein. Representative examples of recombinant pox viruses include ALVAC, TROVAC, NYVAC, and vCP205 (ALVAC-MN120TMG). These viruses were deposited under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Md., 20852, USA: NYVAC under ATCC accession number VR-2559 on Mar. 6, 1997; vCP205 (ALVAC-MN120TMG) under ATCC accession number VR-2557 on Mar. 6, 1997;

TROVAC under ATCC accession number VR-2553 on Feb. 6, 1997 and, ALVAC under ATCC accession number VR-2547 on Nov. 14, 1996.

NYVAC is a genetically engineered vaccinia virus strain generated by the specific deletion of eighteen open reading frames encoding gene products associated with virulence and host range. NYVAC is highly attenuated by a number of criteria including: i) decreased virulence after intracerebral inoculation in newborn mice, ii) inocuity in genetically ( $nu^+/nu^+$ ) or chemically (cyclophosphamide) immunocompromised mice, iii) failure to cause disseminated infection in immunocompromised mice, iv) lack of significant induration and ulceration on rabbit skin, v) rapid clearance from the site of inoculation, and vi) greatly reduced replication competency on a number of tissue culture cell lines including those of human origin.

TROVAC refers to an attenuated fowlpox that was a plaque-cloned isolate derived from the FP-1 vaccine strain of fowlpoxvirus which is licensed for vaccination of 1 day old chicks.

ALVAC is an attenuated canarypox virus-based vector that was a plaque-cloned derivative of the licensed canarypox vaccine, Kanapox (Tartaglia *et al.*, *AIDS Res Hum Retroviruses* 8:1445-7 (1992)). ALVAC has some general properties which are the same as some general properties of Kanapox. ALVAC-based recombinant viruses expressing extrinsic immunogens have also been demonstrated efficacious as vaccine vectors. This avipox vector is restricted to avian species for productive replication. In human cell cultures, canarypox virus replication is aborted early in the viral replication cycle prior to viral DNA synthesis. Nevertheless, when engineered to express extrinsic immunogens, authentic expression and processing is observed *in vitro* in mammalian cells and inoculation into numerous mammalian species induces antibody and cellular immune responses to the extrinsic immunogen and provides protection against challenge with the cognate pathogen.

NYVAC, ALVAC and TROVAC have also been recognized as unique among all poxviruses in that the National Institutes of Health ("NIH")(U.S. Public Health Service), Recombinant DNA Advisory Committee, which issues guidelines for the physical containment of genetic material such as viruses and vectors, *i.e.*, guidelines for safety procedures for the use of such viruses and vectors which are based upon the pathogenicity of the particular virus or vector, granted a reduction in physical containment level: from BSL2 to BSL1. No other poxvirus has a BSL1 physical

containment level. Even the Copenhagen strain of vaccinia virus-the common smallpox vaccine-has a higher physical containment level; namely, BSL2. Accordingly, the art has recognized that NYVAC, ALVAC and TROVAC have a lower pathogenicity than any other poxvirus.

5 Another attenuated poxvirus of preferred use for this invention is Modified Vaccinia virus Ankara (MVA), which acquired defects in its replication ability in humans, as well as most mammalian cells, following over 500 serial passages in chicken fibroblasts (*see, e.g., Mayr et al., Infection* 3:6-14 (1975); Carrol, M. and Moss, B. *Virology* 238:198-211 (1997)). MVA retains its original immunogenicity  
10 and its variola-protective effect and no longer has any virulence and contagiousness for animals and humans. As in the case of NYVAC or ALVAC, expression of recombinant protein occurs during an abortive infection of human cells, thus providing a safe, yet effective, delivery system for foreign antigens.

The HIV antigen encoding DNA for insertion into these vectors is any that is  
15 known to be an effective antigen for protection against a retrovirus. For HIV these would include nucleic acid that can encode at least one of: HIV1 gag( + pro)(IIIB), gp120(MN)( + transmembrane), nef(BRU)CTL, pol(IIIB)CTL, ELDKWA or LDKW epitopes, preferably HIV1 gag( + pro)(IIIB), gp120(MN) ( + transmembrane), two (2) nef(BRU)CTL and three (3) pol(IIIB)CTL epitopes; or two ELDKWA in gp120 V3 or  
20 another region or in gp160. The two (2) nef(BRU)CTL and three (3) pol(IIIB)CTL epitopes are preferably CTL1, CTL2, pol1, pol2 and pol3. In the above listing, the viral strains from which the antigens are derived are noted parenthetically.

#### *Direct DNA delivery vaccines*

25 As an alternative to a viral vaccine, the nucleic acid can also be directly introduced into the cells of a patient. This approach is described, for instance, in Wolff *et. al., Science* 247:1465 (1990) as well as U.S. Patent Nos. 5,580,859; 5,589,466; 5,804,566; 5,739,118; 5,736,524; 5,679,647; and WO 98/04720. Examples of DNA-based delivery technologies include, "naked DNA", facilitated  
30 (bupivacaine, polymers, peptide-mediated) delivery, and cationic lipid complexes or liposomes. The nucleic acids can be administered using ballistic delivery as described, for instance, in Fynan *et al., Proc Natl Acad Sci U S A.* 90:11478-82 (1993) and U.S. Patent No. 5,204,253 or pressure (*see, e.g., U.S. Patent No. 5,922,687*). Using this technique, particles comprised solely of DNA are administered, or in an

alternative embodiment, the DNA can be adhered to particles, such as gold particles, for administration.

As is well known in the art, a large number of factors can influence the efficiency of expression of antigen genes and/or the immunogenicity of DNA vaccines. Examples of such factors include the reproducibility of inoculation, construction of the plasmid vector, choice of the promoter used to drive antigen gene expression and stability of the inserted gene in the plasmid.

Any of the conventional vectors used for expression in eukaryotic cells may be used for directly introducing DNA into tissue. Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, *e.g.*, SV40 vectors. Other exemplary eukaryotic vectors include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of such promoters as the SV40 early promoter, SV40 later promoter, metallothionein promoter, human cytomegalovirus promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

Therapeutic quantities of plasmid DNA can be produced for example, by fermentation in *E. coli*, followed by purification. Aliquots from the working cell bank are used to inoculate growth medium, and grown to saturation in shaker flasks or a bioreactor according to well known techniques. Plasmid DNA can be purified using standard bioseparation technologies such as solid phase anion-exchange resins. If required, supercoiled DNA can be isolated from the open circular and linear forms using gel electrophoresis or other methods.

Purified plasmid DNA can be prepared for injection using a variety of formulations. The simplest of these is reconstitution of lyophilized DNA in sterile phosphate-buffer saline (PBS). This formulation, known as "naked DNA," is particularly suitable for intramuscular (IM) or intradermal (ID) administration.

To maximize the immunotherapeutic effects of plasmid DNA vaccines, alternative methods for formulating purified plasmid DNA may be desirable. A variety of methods have been described, and new techniques may become available. Cationic lipids can also be used in the formulation (*see, e.g.*, as described by WO 93/24640; Mannino & Gould-Fogerite, *BioTechniques* 6(7): 682 (1988); U.S. Pat No. 5,279,833; WO 91/06309; and Felgner, *et al.*, *Proc. Nat'l Acad. Sci. USA* 84:7413

(1987). In addition, glycolipids, fusogenic liposomes, peptides and compounds referred to collectively as protective, interactive, non-condensing compounds (PINC) could also be complexed to purified plasmid DNA to influence variables such as stability, intramuscular dispersion, or trafficking to specific organs or cell types.

5

#### **Selection of an HIV specific epitope.**

Highly antigenic epitopes for provoking an immune response selective for a specific retroviral pathogen are known. The retrovirus, HIV is a major problem in the United States and in the world. With minor exceptions, the following discussion of HIV epitopes is applicable to other retroviruses except for the differences in sizes of the respective viral proteins. HIV-specific epitopes fall into two major categories, structural and non-structural proteins. Epitopes can be selected from either or both groups of proteins. Structural proteins are a physical part of the virion. Non-structural proteins are regulatory proteins. The envelope is a preferred source of epitopes and gp160, 120 and 41 are sources of immunoprotective proteins. Both B and T cell epitopes have been described in the literature and can be used. Peptides selected from the V3 loop of the HIV envelope proteins are of preferred use. In addition other structural proteins have been reported to be immunoprotective including p41, p17 and the gag protein. Non-structural genes include the *rev*, *tat*, *nef*, *vif*, and *vpr* genes.

20

#### **Patients**

A preferred patient population of retrovirally infected persons are those that exhibit repeated and sustained proliferative T-cell responses to envelope epitopes, *e.g.*, HIV gp120. More preferred are those patients that also respond to the *gag* epitopes, *e.g.* HIV p24. Typically these patients are identified by measuring the ability of their lymphocytes to proliferate in responses to highly purified antigen. In brief, peripheral blood monocytes (PBMC) are collected and cultured in the absence of IL-2 and in the presence of 10 µg of highly purified antigen. After four days the cultures are harvested and proliferation is measured by uptake of radioactive thymidine.

30

An alternative means of identifying these patients is to use a skin test. Skin tests involve the detection of a delayed type hypersensitive response (DTH) by means of injecting or scratching antigen beneath the surface of the skin. The reaction is measured by the ability or inability of a patient to exhibit hypersensitive response to an aqueous solution of a gp120

or p24 antigen. Approximately, 1-20  $\mu$ g is applied. The reaction is determined by measuring wheal sizes from about 24 to about 72 hours after administration of a sample, and more preferably from about 48 hours to about 72 hours after administration of a sample. Preferred wheal sizes for evaluation of the hypersensitivity of an animal range from about 16 mm to about 8 mm, more preferably from about 15 mm to about 9 mm, and even more preferably from about 14 mm to about 10 mm in diameter.

#### *Highly Active Anti-Retroviral Therapy (HAART)*

Antiviral retroviral treatment involves the use of two broad categories of therapeutics. They are reverse transcriptase inhibitors and protease inhibitors. There are two type of reverse transcriptase inhibitors: nucleoside analog reverse transcriptase inhibitors and non-nucleoside reverse transcriptase inhibitors. Both types of inhibitors block infection by blocking the activity of the HIV reverse transcriptase, the viral enzyme that translates HIV RNA into DNA which can later be incorporated into the host cell chromosomes.

Nucleoside and nucleotide analogs mimic natural nucleotides, molecules that act as the building blocks of DNA and RNA. Both nucleoside and nucleotide analogs must undergo phosphorylation by cellular enzymes to become active; however, a nucleotide analog is already partially phosphorylated and is one step closer to activation when it enters a cell. Following phosphorylation, the compounds compete with the natural nucleotides for incorporation by HIV's reverse transcriptase enzyme into newly synthesized viral DNA chains, resulting in chain termination.

Examples of anti-retroviral nucleoside analogs are: AZT, ddI, ddC, d4T, and 3TC in combination with AZT and Combivir.

Nonnucleoside reverse transcriptase inhibitors (NNRTIs) are a structurally and chemically dissimilar group of antiretroviral compounds. They are highly selective inhibitors of HIV-1 reverse transcriptase. At present these compounds do not affect other retroviral reverse transcriptase enzymes such as hepatitis viruses, herpes viruses, HIV-2, and mammalian enzyme systems. They are used effectively in triple-therapy regimes. Examples of NNRTIs are Delavirdine and Nevirapine which have been approved for clinical use in combination with nucleoside analogs for treatment of HIV-infected adults who experience clinical or immunologic deterioration. A detailed review can be found in "Nonnucleoside Reverse Transcriptase Inhibitors" *AIDS Clinical Care* (10/97) Vol. 9, No. 10, p. 75.

Proteases inhibitors are compositions that inhibit HIV protease, which is virally encoded and necessary for the infection process to proceed. Clinicians in the United States have a number of clinically effective proteases to use for treating HIV-infected persons. These include: SAQUINAVIR (Invirase); INDINAVIR (Crixivan);  
5 and RITONAVIR (Norvir).

#### *CD4<sup>+</sup> T cell counts*

To assess a patient's immune system before antiviral treatment and after treatment as well as to determine if the claimed vaccine regimen is working it is  
10 important to measure CD4<sup>+</sup> T cell counts. A detailed description of this procedure was published by Janet K.A. Nicholson, Ph.D *et al.* 1997 *Revised Guidelines for Performing CD4<sup>+</sup> T-Cell Determinations in Persons Infected with Human Immunodeficiency Virus (HIV)* in *The Morbidity and Mortality Weekly Report*, 46(RR-2):[inclusive page numbers], Feb 14, 1997, Centers for Disease Control.

15 In brief, most laboratories measure absolute CD4<sup>+</sup> T-cell levels in whole blood by a multi-platform, three-stage process. The CD4<sup>+</sup> T-cell number is the product of three laboratory techniques: the white blood cell (WBC) count; the percentage of WBCs that are lymphocytes (differential); and the percentage of lymphocytes that are CD4<sup>+</sup> T-cells. The last stage in the process of measuring the percentage of CD4<sup>+</sup> T-  
20 lymphocytes in the whole-blood sample is referred to as "immunophenotyping by flow cytometry.

Immunophenotyping refers to the detection of antigenic determinants (which are unique to particular cell types) on the surface of WBCs using antigen-specific monoclonal antibodies that have been labeled with a fluorescent dye or fluorochrome  
25 (e.g., phycoerythrin [PE] or fluorescein isothiocyanate [FITC]). The fluorochrome-labeled cells are analyzed by using a flow cytometer, which categorizes individual cells according to size, granularity, fluorochrome, and intensity of fluorescence. Size and granularity, detected by light scattering, characterize the types of WBCs (*i.e.*, granulocytes, monocytes, and lymphocytes). Fluorochrome-labeled antibodies  
30 distinguish populations and subpopulations of WBCs.

Systems for measuring CD4<sup>+</sup> cells are commercially available. For example Becton Dickinson's FACSCount System automatically measure absolutes CD4<sup>+</sup>, CD8<sup>+</sup>, and CD3<sup>+</sup> T lymphocytes. It is a self-contained system, incorporating instrument, reagents, and controls.

### *Viral titer*

There are a variety of ways to measure viral titer in a patient. A review of the state of this art can be found in the *Report of the NIH To Define Principles of Therapy of HIV Infection* as published in the *Morbidity and Mortality Weekly Reports*, April 24, 1998, Vol 47, No. RR-5, Revised 6/17/98. It is known that HIV replication rates in infected persons can be accurately gauged by measurement of plasma HIV concentrations.

HIV RNA in plasma is contained within circulating virus particles or virions, with each virion containing two copies of HIV genomic RNA. Plasma HIV RNA concentrations can be quantified by either target amplification methods (e.g., quantitative RT polymerase chain reaction [RT-PCR], Amplicor HIV Monitor assay, Roche Molecular Systems; or nucleic acid sequence-based amplification, [NASBA<sup>®</sup>], NucliSens<sup>™</sup> HIV-1 QT assay, Organon Teknika) or signal amplification methods (e.g., branched DNA [bDNA], Quantiplex<sup>™</sup> HIV RNA bDNA assay, Chiron Diagnostics). The bDNA signal amplification method amplifies the signal obtained from a captured HIV RNA target by using sequential oligonucleotide hybridization steps, whereas the RT-PCR and NASBA<sup>®</sup> assays use enzymatic methods to amplify the target HIV RNA into measurable amounts of nucleic acid product. Target HIV RNA sequences are quantitated by comparison with internal or external reference standards, depending upon the assay used.

### *Measurements of CD8<sup>+</sup> Responses*

CD8<sup>+</sup> T-cell responses can be measured, for example, by using tetramer staining of fresh or cultured PBMC (see, e.g., Altman, J. D. *et al.*, *Proc. Natl. Acad. Sci. USA* 90:10330, 1993; Altman, J. D. *et al.*, *Science* 274:94, 1996), or  $\gamma$ -interferon release assays such as ELISPOT assays (see, e.g., Lalvani, A. *et al.*, *J. Exp. Med.* 186:859, 1997; Dunbar, P. R. *et al.*, *Curr. Biol.* 8:413, 1998; Murali-Krishna, K. *et al.*, *Immunity* 8:177, 1998), or by using functional cytotoxicity assays. Each of these assays are well-known to those of skill in the art. For example, a cytotoxicity assay can be performed as follows.

Briefly, peripheral blood lymphocytes from patients are cultured with HIV peptide epitope at a density of about five million cells/ml. Following three days of culture, the medium is supplemented with human IL-2 at 20 units/ml and the cultures

are maintained for four additional days. PBLs are centrifuged over Ficoll-Hypaque and assessed as effector cells in a standard  $^{51}\text{Cr}$ -release assay using U-bottomed microtiter plates containing about  $10^4$  target cells with varying effector cell concentrations. All cells are assayed twice. Autologous B lymphoblastoid cell lines  
5 are used as target cells and are loaded with peptide by incubation overnight during  $^{51}\text{Cr}$  labeling. Specific release is calculated in the following manner: (experimental release-spontaneous release)/(maximum release-spontaneous release) x 100. Spontaneous release is generally less than 20% of maximal release with detergent (2% Triton X-100) in all assays.

10

### Formulation of Vaccines and Administration

The administration procedure for recombinant virus or DNA is not critical. Vaccine compositions (*e.g.*, compositions containing the poxvirus recombinants or DNA) can be formulated in accordance with standard techniques well known to those  
15 skilled in the pharmaceutical art. Such compositions can be administered in dosages and by techniques well known to those skilled in the medical arts taking into consideration such factors as the age, sex, weight, and condition of the particular patient, and the route of administration.

For example, NYVAC-HIV or other attenuated pox virus vaccines such as  
20 ALVAC-HIV or MVA-HIV, is inoculated, more than once, by the intramuscular route at a dose of about  $10^8$  pfu per inoculation, for a patient of 170 pounds. The vaccine can be delivered in a physiologically compatible solution such as sterile PBS in a volume of, *e.g.*, one ml. The dose can be proportional to weight.

The compositions can be administered alone, or can be co-administered or  
25 sequentially administered with other immunological, antigenic, vaccine, or therapeutic compositions. Such compositions can include other agents to potentiate or broaden the immune response, *e.g.*, IL-2 or CD40 ligand, which can be administered at specified intervals of time, or continuously administered (*see, e.g.*, Smith *et al.*, *N Engl J Med* 1997 Apr 24;336(17):1260-1; and Smith, *Cancer J Sci Am.* 1997 Dec;3  
30 Suppl 1:S137-40). For example, IL-2 can be administered in a broad range, *e.g.*, from 10,000 to 1,000,000 or more units. Administration can occur continuously following vaccination. Often, low doses, *e.g.* 100,000 to 200,000, often 120,000, 150,000 or 170,000, units of IL-2 can be particularly useful.

Other compositions that can be co-administered can include purified antigens from immunodeficiency virus or antigens that are expressed by a second recombinant vector system which is able to produce other therapeutic compositions. Such compositions can include a recombinant poxvirus which expresses other immunodeficiency antigens or biological response modifiers (*e.g.* cytokines; co-stimulating molecules). Again, co-administration is performed by taking into consideration such known factors as the age, sex, weight, and condition of the particular patient, and, the route of administration.

DNA expression vectors for direct introduction of DNA into the patient tissue can additionally be complexed with other components such as peptides, polypeptides and carbohydrates. Expression vectors can also be complexed to particles or beads that can be administered to an individual, for example, using a vaccine gun.

The expression vectors are administered by methods well known in the art as described in Donnelly *et al.* (*Ann. Rev. Immunol.* 15:617-648 (1997)); Felgner *et al.* (U.S. Patent No. 5,580,859, issued December 3, 1996); Felgner (U.S. Patent No. 5,703,055, issued December 30, 1997); and Carson *et al.* (U.S. Patent No. 5,679,647, issued October 21, 1997), each of which is incorporated herein by reference. One skilled in the art would know that the choice of a pharmaceutically acceptable carrier, including a physiologically acceptable compound, depends, for example, on the route of administration of the expression vector.

For example, naked DNA or polynucleotide in an aqueous carrier can be injected into tissue, such as muscle, in amounts of from 10  $\mu$ l per site to about 1 ml per site. The concentration of polynucleotide in the formulation is from about 0.1  $\mu$ g/ml to about 20 mg/ml.

Vaccines can be delivered via a variety of routes. Typical delivery routes include parenteral administration, *e.g.*, intradermal, intramuscular or subcutaneous routes. Other routes include oral administration, intranasal, and intravaginal routes.

The expression vectors of use for the invention can be delivered to the interstitial spaces of tissues of a patient (*see, e.g.*, Felgner *et al.*, U.S. Patent Nos. 5,580,859, and 5,703,055). Administration of expression vectors of the invention to muscle is a particularly effective method of administration, including intradermal and subcutaneous injections and transdermal administration. Transdermal administration, such as by iontophoresis, is also an effective method to deliver expression vectors of the invention to muscle. Epidermal administration of expression vectors of the invention can also be employed. Epidermal

administration involves mechanically or chemically irritating the outermost layer of epidermis to stimulate an immune response to the irritant (Carson *et al.*, U.S. Patent No. 5,679,647).

The vaccines can also be formulated for administration via the nasal passages.

5 Formulations suitable for nasal administration, wherein the carrier is a solid, include a coarse powder having a particle size, for example, in the range of about 10 to about 500 microns which is administered in the manner in which snuff is taken, *i.e.*, by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid for administration  
10 as, for example, nasal spray, nasal drops, or by aerosol administration by nebulizer, include aqueous or oily solutions of the active ingredient. For further discussions of nasal administration of AIDS-related vaccines, references are made to the following patents, US 5,846,978, 5,663,169, 5,578,597, 5,502,060, 5,476,874, 5,413,999, 5,308,854, 5,192,668, and 5,187,074.

15 Examples of vaccine compositions of use for the invention include liquid preparations, for orifice, *e.g.*, oral, nasal, anal, vaginal, etc. administration, such as suspensions, syrups or elixirs; and, preparations for parenteral, subcutaneous, intradermal, intramuscular or intravenous administration (*e.g.*, injectable administration) such as sterile suspensions or emulsions. In such compositions the  
20 recombinant poxvirus, expression product, immunogen, DNA, or modified gp120 or gp160 can be in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose or the like.

The vaccines can be incorporated, if desired, into liposomes, microspheres or other polymer matrices (*see, e.g.*, Felgner *et al.*, U.S. Patent No. 5,703,055;  
25 Gregoriadis, *Liposome Technology*, Vols. I to III (2nd ed. 1993). Liposomes, for example, which consist of phospholipids or other lipids, are nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer. Liposomes include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like.

30 Liposome carriers can serve to target a particular tissue or infected cells, as well as increase the half-life of the vaccine. In these preparations the vaccine to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to, *e.g.*, a receptor prevalent among lymphoid cells, such as monoclonal antibodies which bind to the CD45 antigen, or with other therapeutic or immunogenic compositions. Thus, liposomes

either filled or decorated with a desired immunogen of the invention can be directed to the site of lymphoid cells, where the liposomes then deliver the immunogen(s).

Liposomes for use in the invention are formed from standard vesicle-forming lipids, which generally include neutral and negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, *e.g.*, liposome size, acid lability and stability of the liposomes in the blood stream. A variety of methods are available for preparing liposomes, as described in, *e.g.*, Szoka, *et al.*, *Ann. Rev. Biophys. Bioeng.* 9:467 (1980), U.S. Patent Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

## EXAMPLES

The following examples are provided by way of illustration only and not by way of limitation. Those of skill will readily recognize a variety of noncritical parameters which could be changed or modified to yield essentially similar results.

### **Example 1. Administration of NYVAC-SIV<sub>gag-pol-env</sub> to SIV-infected, HAART-treated *Rhesus macaques***

The effectiveness of a highly attenuated Poxvirus vector as a therapeutic vaccine to enhance host-immune responses was investigated in the SIV<sub>251</sub> Rhesus macaque model, which models HIV-1 infection in humans. The vaccine used for this study was a highly attenuated NYVAC-SIV<sub>gag-pol-env</sub> recombinant vaccine that was demonstrated to have efficacy as a preventative vaccine in earlier studies (Benson *et al.*, *J. Virol.* 72:4170-4182 (1998)).

The study design included 24 animals which were divided into three groups, A, B, and C. All the animals were infected intravenously with ten infectious doses of highly pathogenic SIV<sub>251</sub> (Pal). Following SIV<sub>251</sub> exposure, all twenty four

animals became infected; the peak of plasma viremia occurred at approximately two weeks and ranged between  $10^7$  and  $10^9$  copies of viral RNA/ml of plasma in the twenty four animals.

Two and a half weeks after infection, sixteen animals, those in groups A and B, received a HAART regimen that in a pilot study had reduced viremia to undetectable levels in 80% of macaques chronically infected with SIV<sub>251</sub>. The HAART regimen included oral administration of two doses of Stavudine (1.2 mg/day), intravenous inoculation of DDI (10 mg/kg/day) and subcutaneous inoculation of PMPA (20 mg/kg/day). Animals in group C were not treated with drugs. Animals in groups A and B, but not C, experienced a significant decrease in viremia (Figure 1, top panels).

In the sixteen animals of groups A and B, HAART treatment was continued daily for 6 months. At weeks 10, 19, and 23 post-infection, the animals in group A received a placebo vaccine (non-recombinant NYVAC vector) and the animals in groups B and C received  $10^8$  pfu of NYVAC-SIV *gag-pol-env* vaccine. All twenty-four animals were monitored weekly for viral RNA copies/ml of plasma and biweekly for lymphoproliferative responses (LPR) to highly purified native p27 *gag* and gp120 *env* SIV proteins.

In order to follow the CD8<sup>+</sup> T-cell responses, three MAMU-A\*01 (*Macaca mulata* equivalent of HLA class I A\*01 (Kuroda *et al.*, *J. Exp. Med.* 187:1373-1381, 1998) were included in each group. MAMU-A\*01 animals are generally able to recognize the immunodominant peptide 11c-m within the gag antigen of SIV. A tetramer binding assay was therefore used to directly quantitate CD8<sup>+</sup> T-cell responses *in vivo*. The tetramer, formed by four identical MAMU-A\*01 molecules conjugated to the peptide 11c-m, was linked to fluorescent-labeled streptavidin and used to stain the CD8<sup>+</sup> T-cells in the blood of macaques that expressed the appropriate T cell receptor complex on their surface. The percentage of total CD8<sup>+</sup>/CD3<sup>+</sup> staining with the peptide MAMU-A\*-1 tetramer was measured in several consecutive time intervals in each of the MAMU-A\*01 animals.

Staining of fresh or cultured PBMC with the MAMU A\*01/peptide 11c-m tetramer was performed in all the nine MAMU A\*01 animals included in the study (three in each group) following SIV<sub>251</sub> infection and after the second NYVAC-SIV vaccination. MAMU-A\*01/peptide-11c-m-tetramer-staining CD8<sup>+</sup> T-cells (ranging from 0.8 to 4.6%) were induced by SIV infection in all nine animal within the first

month after viral exposure, and the CD8<sup>+</sup> T-cell population was expanded (up to values of approximately 70%) *in vitro* following specific peptide 11c-m stimulation. At two months after SIV<sub>251</sub> exposure, the infected macaques seroconverted to SIV<sub>251</sub> antigens, including the animals in groups A and B that were treated with HAART.

5 Measurements of LPR to gp120 and p27 *gag* was consistently negative in all twenty four animals within the first four weeks following viral infection.

*The frequency and extent of CD4<sup>+</sup> T-helper responses in HAART-treated animals is increased by NYVAC-SIV vaccination.*

10 As stated above, acute infection by SIV<sub>251</sub> was associated with the absence of a proliferative response to both p27 *gag* and gp120 *env*. However, responses to p27 *gag* appeared in HAART-treated animals at approximately 10 weeks after infection and, these responses were more frequent in animals of group A than in animals of group C. No difference in the LPR to gp120 was observed between these two groups  
15 (Figure 1, middle and lower panels).

This notion was further supported by the finding that two animals (647 and 655) in group B failed to respond to therapy, maintained high virus load, and did not develop CD4<sup>+</sup> T-cell proliferative response following NYVAC-SIV vaccination.

Further corroborating this notion, two animals in group C that naturally  
20 controlled viremia developed LPR following NYVAC-SIV vaccination. Thus, it appears that CD4<sup>+</sup> T-helper memory responses are induced inefficiently by NYVAC-SIV animals with high viremia. Several factors may contribute to this finding: CD4<sup>+</sup> T-cells are already activated *in vivo* and do not further proliferate *in vitro* in LPR assays and/or the vaccine-induced memory cells become targets for SIV infection and  
25 die upon further antigen stimulation. These data provide the first evidence that a highly attenuated live recombinant poxvirus vector vaccine can induce and boost sustained CD4<sup>+</sup> helper immune response in the context of a pharmacologically controlled lentiviral infection.

30 *NYVAC-SIV vaccination increases CD8<sup>+</sup>/CD3<sup>+</sup> MAMU-A\*01-tetramer-positive cells only in HAART-treated animals.*

SIV infection induced a large number of CD8<sup>+</sup> T-cells that bound the MAMU-A\*01 tetramer. This response by week four was reduced in most animals. Following the second and third NYVAC-SIV vaccinations, a high percentage of CD8<sup>+</sup>/CD3<sup>+</sup> T-

cells bound tetramers in the fresh PBMC of all MAMU-A\*01 animals in group B, but in none of the animals in groups C (Figure 2). The specificity of tetramer staining was shown in parallel experiments using PBMC from animals with a different haplotype as well as by the expansion of the cells from the nine MAMU-A\*01 animals *in vitro* following peptide-11c-m-specific stimulation. Peptide 11c-m presented in the context of the MAMU-A\*01 haplotype is recognized by CD8<sup>+</sup> T-cells with cytolytic activity. (In Figure 2, the top panel shows the results obtained in the three MAMU-A\*01 animals from group A; middle panel, from group B; and bottom panel, from group C. The percentage of MAMU-A\*01/peptide-11c-m-tetramer-staining cells within the first 4 weeks was evaluated using only α-CD8<sup>+</sup> antibodies as a T cell marker whereas the data presented from weeks 19 through 29 were obtained using simultaneously α-CD-3<sup>+</sup> and α-CD8<sup>+</sup> antibodies in conjunction with the MAMU-A\*01/peptide-11c-m tetramers. For some of the MAMU-A\*01-positive animals, CTL activity obtained in cultured PBMC [weeks 19 and 20] or fresh PBMC [week 23] is also presented. The numbers in the abscissa represent the effector-target ratio of the CTL assay system.)

To assess whether the detection of MAMU-A\*01/peptide-11c-m-tetramer-staining CD8<sup>+</sup>/CD3<sup>+</sup> T-cells in the PBL of macaques was associated with CTL activity, cytotoxicity assays were performed using homologous B cells from each animal pulsed with peptide 11 c-m. It was shown that CTL activity was measured after *in vitro* stimulation of CD8<sup>+</sup> T-cells in all animals tested at weeks 19 and 20, although the extent of killing did not correlate with the percentage of tetramer-staining cells (Figure 2). Most notably, CTL assays performed on fresh CD8<sup>+</sup> T cells at day 23 demonstrated significant CTL activity in group B animals, confirming the CTL functional activity of the high number of circulating CD8<sup>+</sup> tetramer-staining cells in the blood of these animals (Figure 2). Thus, NYVAC-SIV vaccination induced high levels of CD8<sup>+</sup> responses only in animals in which viral replication was suppressed by therapy.

#### *Delayed T-cell hypersensitivity (DTH) to viral p27 gag*

Vaccines able to induce T-cell-mediated immunity are often able to induce DTH. To assess whether any of the animals vaccinated with NYVAC-SIV developed this response, either 1 or 10 ug of highly purified SIV p27 or HTLV-I p24, as controls, were inoculated

intradermally in animals in groups B and C. DTH reactivity was considered positive when a thickness of more than 10 mm manifested at 72 h postinoculation. Only three animals in group B and two animals in group C fulfilled the requirement for DTH positivity.

The data presented in this example demonstrated that inoculations of NYVAC-SIV following HAART greatly increased the frequency, extent, and duration of those responses in animals in which viremia was efficiently suppressed, indicating that the ability to detect vaccine-induced CD4<sup>+</sup> T-cell helper responses was strictly dependent on the level of viral replication in the host. Similarly, NYVAC-SIV vaccination induced significant expansion of the number of CD8<sup>+</sup>/CD3<sup>+</sup> MAMU-A\*01 cells specific for an immunodominant SIV *gag* peptide only in animals treated effectively with antiviral therapy. Following therapy suspension, NYVAC-SIV-vaccinated animals were able to control viremia better than animals treated with antiviral therapy alone. These data demonstrate that vaccination can further induce both CD4<sup>+</sup> and CD8<sup>+</sup> T-cell responses in SIV-infected macaques.

## **Example 2. Immunization of a person infected with HIV.**

A 35 year old male patient is seropositive for HIV and has a viral count in his plasma of 15000 copies per milliliter and a CD4<sup>+</sup> count of 300. The patient is treated with a cocktail of antiviral agents which consist of: two nucleotide inhibitors and one protease inhibitor (Zaduvudine [three time a day], Lamivudine [two times a day], and Nesinavir [three times day] at preformulated doses. Typically, antiretroviral therapy is considered effective if a decrease of at least one log of the plasma viral RNA is observed. The plasma RNA load as well as the CD4<sup>+</sup> T-cell count in the blood will be measured monthly.

After six months the patient is re-evaluated and is determined to have a viral count in his plasma of 1500 copies per milliliter or less and a CD4<sup>+</sup> count of 500. He is then injected with an attenuated pox virus vector NYVAC carrying the following viral peptides: *gag*, *pro*, *gp120-TM*, *pol* and *nef* string of CTL epitopes. The injection comprises 10<sup>8</sup> pfu of the pox virus.

The patient's immune responses is evaluated (CD4+ proliferative response; cytotoxic CD8+ T-cell activity, etc.) and a decision is made as to whether and when to immunize again. Typically, a maximum of three to four immunizations with NYVAC-HIV-1 is considered. This regimen could be followed by three to four immunizations with ALVAC-HIV-1 (carrying a similar HIV-I genetic content) and, if

necessary, an additional regimen of DNA-only *i.e.*, DNA that is not in a viral vector, immunization could follow. NYVAC-HIV-1 followed by ALVAC-HIV-1 is effective in inducing immunoresponses in chimpanzees. Similarly, immunization regimens as follows have been shown to be effective: NYVAC followed by DNA and ALVAC followed by DNA. Thus, a vaccination regimen including all of the above vaccines may be used.

Often, the vaccine regimen is administered with IL-2, preferably at low doses such as 100,000 to 200,000 units of IL-2 administered daily. CD40<sup>+</sup> ligand can also be included in the treatment protocol, either by itself or administered in conjunction with the IL-2 treatment.

**Example 3. Administration of ALVAC-SIV<sub>gpc</sub> to SIV-infected, HAART-treated *Rhesus macaques***

The effectiveness of a highly attenuated ALVAC vector as a therapeutic vaccine to enhance host-immune responses in HAART-treated animals was investigated in the SIV<sub>251</sub> *Rhesus macaque* model, which models HIV-1 infection in humans. The study includes 16 macaques inoculated with the SIV<sub>251</sub> virus and treated with the HAART regimen as described in Example 1 at day 15 and thereafter. Of those macaques, 8 are immunized with a total of 3 doses of 10<sup>8</sup> pfu of the mock-vaccine ALVAC vector (group D) and the remaining 8 (group E) with a recombinant ALVAC-SIV-*gag-pol-env* vector (ALVAC-SIV<sub>gpc</sub>), which is analogous to the NYVAC vector of Example 1.

The data obtained after a single dose of the ALVAC-SIV<sub>gpc</sub> vaccine indicated that the ALVAC-SIV<sub>gpc</sub> vaccine is able to boost CD4<sup>+</sup> T-cell responses against the p27 Gag protein as well as to the gp120 Env forefront of the vaccine (see Figure 3). In addition, a specific CD8<sup>+</sup> T-cell response detected using the MAMU A\*01/peptide 11c-m tetramer reagent (*see, e.g.*, Example 1) was also boosted by the ALVAC-SIV<sub>gpc</sub> (Figure 4), and further, those CD8<sup>+</sup> T-cells could be expanded in culture. The mock-vaccinated animals did not experience an expansion of these immune responses.

Thus, ALVAC-SIV<sub>gpc</sub> is immunogenic in animals undergoing HAART therapy. An ALVAC vaccine analogous to the NYVAC vaccine of Example 2 can similarly be employed to treat HIV-infected individuals as described in Example 2 who are undergoing HAART.

**Example 4. MVA-SIV-*gag-pol-env* immunization of SIV-infected macaques.**

The effectiveness of an MVA-SIV vaccine was evaluated using methodology analogous to that used to evaluate NYVAC and ALVAC SIV *gag-pol-env* vaccines. The CD8<sup>+</sup> and CD4<sup>+</sup> responses were determined in SIV<sub>251</sub>-infected macaques that were able to control viremia, *i.e.*, the CD4<sup>+</sup> T-cell counts were above 500, following administrations of a single dose of 10<sup>8</sup> pfu of MVA-SIV-*gag-pol-env* recombinant vaccine. The results showed that both CD4<sup>+</sup> and CD8<sup>+</sup> responses could be expanded in infected animals.

Thus, MVA-SIV-*gag-pol-env* is also immunogenic in animals and can be used in patients undergoing HAART therapy.

**Example 5. Comparison of the Immunogenicity of DNA and NYVAC vaccines.**

A vaccine regimen comprising administration of DNA alone or DNA in combination with NYVAC-SIV or ALVAC-SIV, or administration of a combination of NYVAC-SIV and ALVAC-SIV is also effective in continuous boost of the immune response in SIV<sub>251</sub>-infected animals.

A study conducted in parallel compared 2 inoculations of 10<sup>8</sup> pfu of NYVAC-SIV<sub>gpc</sub> and 3 inoculations of DNA, which was administered using 4mg intramuscularly and 1 mg intradermally of each plasmid, in naïve animals. The DNA vaccine induced CD4<sup>+</sup> and CD8<sup>+</sup> T-cell responses that were equivalent to those induced by a NYVAC-SIV<sub>gpc</sub> vaccine (Figures 5 and 6). ALVAC-SIV<sub>gpc</sub> was at least as immunogenic as NYVAC-SIV<sub>gpc</sub> (*see, e.g.*, Example 1) and NYVAC-SIV<sub>gpc</sub> was as immunogenic as DNA. Thus, all three vaccines either alone or in various combinations can be used in HIV-I-infected individuals.

ALVAC-SIV<sub>gpc</sub> was also able to induce both CD4<sup>+</sup> and CD8<sup>+</sup> T-cell response in chronically infected animals (CD4<sup>+</sup> T-cell range 50-900) and was able to suppress viremia in the absence of HAART. These animals, which had been previously vaccinated with a NYVAC-SIV<sub>gpc</sub>-based vaccine, appear to model long-term progressor HIV-I-infected individuals. The vaccines of the invention can therefore be used alone or in various interchangeable combinations in early infection as well as late infection in individuals in which viremia is controlled pharmacologically or otherwise.

**Example 6. Therapeutic Vaccination with NYVAC and IL-2.**

This examples demonstrates the ability of immunomodulatory molecules, *e.g.* IL-2, to further expand both CD4<sup>+</sup> and CD8<sup>+</sup> T-cell responses induced

by the NYVAC-SIV<sub>gag-pol-env</sub> vaccine and increase the breadth of the host response to the virus.

SIV<sub>251</sub>-infected HAART-treated Rhesus macaques (Figure 7) were vaccinated intramuscularly with NYVAC-SIV<sub>gag-pol-env</sub> with or without simultaneous and continued daily treatment with IL-2 (120,000 IU) administered subcutaneously. A control group of animals was treated with IL-2 and mock-vaccinated (NYVAC nonrecombinant vector). All 15 macaques in the study responded to HAART (10 mg/kg/day DDI, intravenously; 2.4 mg/kg/day Stavudine, orally; 10 mg/kg/day PMPA, subcutaneously) and in 13 animals viremia was suppressed below  $5 \times 10^3$  copies/ml within the first 4 weeks of treatment. Viremia in the remaining 2 animals became undetectable by weeks 6 and 8 of treatment. Proliferative responses to p27 Gag and gp120 were increased by NYVAC-SIV<sub>gag-pol-env</sub> vaccination up to three- and twelve-fold, respectively, regardless of IL-2 treatment, indicating that either IL-2 does not increase bulk proliferative response or that the assay was not sensitive enough to measure subtle variation in the antigen-specific CD4<sup>+</sup> T-helper response.

Approximately half of the animals in the study were genetically selected as carriers of the Mamu-A\*01 molecules. CTL CD8<sup>+</sup> T-cell responses to SIV<sub>251</sub> were therefore measured by ELISPOT using several purified SIVmac<sub>239</sub> nonamer peptides and their corresponding Mamu-A\*01 tetramers. The *ex vivo* PBMC of all SIV<sub>251</sub>-infected Mamu-A\*01 animals recognized the immunodominant p11C,C→M peptide and produced  $\gamma$ -interferon ( $\gamma$ -INF) following *in vitro* stimulation. This response was further expanded following immunization with NYVAC-SIV<sub>gag-pol-env</sub>. While IL-2 did not expand the number of  $\gamma$ -INF-producing cells in response to the p11c, C→M peptide in mock-vaccinated animals, the expansion of this response was higher in the NYVAC-SIV<sub>gag-pol-env</sub>-treated macaques that also received IL-2 than in those that received NYVAC-SIV<sub>gag-pol-env</sub> alone. In contrast, IL-2 *per se* appeared to expand the immune response to two other immunodominant epitopes within the SIV tat and vif proteins in treated animals. (Vaccination with NYVAC-SIV<sub>gag-pol-env</sub> did not further expand these responses as these antigens are not included in the vaccine.) Moreover, the CD8<sup>+</sup> T-cell responses to 2 subdominant epitopes within the Gag and Env proteins of SIV were clearly expanded following NYVAC-SIV<sub>gag-pol-env</sub> vaccination in macaques that received simultaneous and continuous IL-2 treatment. Thus, the administration of low-dose IL-2 in conjunction with vaccination with the

highly attenuated NYVAC-SIV<sub>gag-pol-env</sub> vaccine potentiated and broadened CD8<sup>+</sup> T-cell functional responses to SIV<sub>251</sub>.

*IL-2 can be used with the vaccine to control viremia after antiretroviral therapy*

5 *interruption*

A HAART-treated macaque was inoculated with NYVAC-SIV<sub>gag-pol-env</sub> at the intervals shown in Figure 8. The macaque also received low dose IL-2, *i.e.*, 120,000 units daily administered subcutaneously. This animal exhibited expanded CD8<sup>+</sup> (top panel, Figure 8) and CD4<sup>+</sup> proliferative (lower panel, Figure 8) responses. Furthermore, as shown in the top panel of Figure 8, transient viral rebound occurred following interruption of HAART treatment and after suspension of the IL2 treatment. Thus, administration of IL-2 in conjunction with vaccination can contribute to the control of viremia after interruption of antiretroviral therapy.

10

## WHAT IS CLAIMED IS:

1. A method of stimulating an efficient CD8<sup>+</sup> response in a human infected with a an HIV or HTLV-1 retrovirus said method comprising:  
5 administering to the human, a nucleic acid-based vaccine, which enters the cells of the human and intracellularly produces HIV- or HTLV-1-specific peptides for presentation on the cell's MHC class I molecules

where said peptides are presented in an amount sufficient to stimulate a protective CD8<sup>+</sup> response and

10 where said human  
i. has a viral load of less than 10,000 viral copies per ml of plasma and a CD4<sup>+</sup> cell count of above 500 cells/ml, and  
ii. has been treated with one or more anti-viral agents, which contributed to a lower viral copy and higher CD4<sup>+</sup> cell count than before treatment.

15 2. A method of claim 1 wherein the human has been treated with anti-viral agents, which resulted in the human having a viral load of less than 1,000 viral copies per ml of blood serum and a CD4<sup>+</sup> cell count of above 500 cells/ml.

20 3. A method of claim 2 wherein the anti-viral agents comprise a combination of protease inhibitors and inhibitors of reverse transcriptase.

4. A method of claim 1 wherein the vaccine is a DNA based vaccine.

25 5. A method of claim 1 wherein the vaccine is an attenuated recombinant virus.

6. A method of claim 5 wherein the vaccine is an attenuated pox virus.

30 7. A method of claim 6 wherein the attenuated pox virus is selected from the group consisting of NYVAC and ALVAC.

8. A method of claim 6 wherein the attenuated pox virus is MVA.

9. A method of claim 1 where the vaccine is administered a second time.

10. A method of claim 1 wherein the HIV- or HTLV-1-specific peptides are structural viral peptides

5

11. A method of claim 1 wherein the HIV- or HTLV-1-specific peptides are non-structural viral peptides.

12. A method of claim 1 wherein the vaccine further comprises an adjuvant.

10

13. A method of claim 1 further comprising administering interleukin 2 or CD40 ligand in an amount sufficient to potentiate the CD8<sup>+</sup> response.

14. A method of claim 1 where the person has been infected with HIV and has demonstrated repeated and sustained proliferative T-cell responses to gp120 envelope protein.

15

15. A method of claim 14 where the person has demonstrated repeated and sustained proliferative T-cell responses to p24 gag antigen.

20

16. A method of claim 1 where the person is infected with HIV and is further tested by a skin test for a hypersensitive response to p24 gag antigen.

17. A method of claim 1 where the person is infected with HIV and is further tested by a skin test for a hypersensitive response to gp120 envelope antigen.

25

# VIRAL LOAD AND CD4<sup>+</sup> T-CELL RESPONSES IN UNTREATED/TREATED NYVAC-SIV VACCINATED MACAQUES




Fig. 1

MAMU-A-01/PEPTIDE 11c TETRAMER STAINING, CD8<sup>+</sup>/CD3<sup>+</sup> T-CELLS -  
AND PEPTIDE 11c SPECIFIC CYTOTOXIC ACTIVITY IN UNTREATED/TREATED  
NYVAC-SIV VACCINATED MACAQUES




Fig. 2

INDUCTION OF PROLIFERATIVE RESPONSE TO p27 gag AND GP120  
BY THE ALVAC-SIV VACCINE IN SIV<sub>251</sub>-INFECTED, HAART-TREATED MACAQUES




Fig. 3

INDUCTION OF CD3<sup>+</sup> CD8<sup>+</sup> TETRAMER-STAINING T-CELLS  
IN SIV<sub>251</sub>-INFECTED, HAART-TREATED MACAQUES  
BY THE CANARYPOX-BASED ALVAC-SIV-*gag-pol-env* VACCINE





Fig. 4a


Fig. 4b

Fig. 4


# INDUCTION OF CD8<sup>+</sup> AND CD4<sup>+</sup> T-CELL RESPONSES BY pCMV-gag AND pCMV-env DNA VACCINE



**Fig. 5b**



**Fig. 5d**



**Fig. 5c**

Fig. 5




Fig. 6

## EXPERIMENTAL DESIGN




Fig. 7




Fig. 8

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/20641

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K39/21

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 08539 A (CHIRON CORPORATION) 5 March 1998 (1998-03-05) the whole document	1-17
X	M. JOHN ET AL.: "Control of HIV replication by cytotoxic T-lymphocyte responses" J. HIV THERAPY, vol. 4, no. 4, 1999, pages 91-87, XP000982254 page 95, column 1, paragraph 3 -page 96	1-17
X	F. GOTCH ET AL.: "Therapeutic Vaccines in HIV 1 infection" IMMUNOLOGICAL REVIEWS, vol. 170, 1999, pages 173-182, XP000982295 page 179, column 2; figure 1	1-17



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

16 March 2001

Date of mailing of the international search report

26/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Skelly, J

# INTERNATIONAL SEARCH REPORT

Int .tional Application No

PCT/US 00/20641

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	R. MACGREGOR ET AL.: "First human trial of a DNA-based vaccine for treatment of HIV1 infection: safety and host response" J. INF. DIS., vol. 178, 1998, pages 92-100, XP000982247 page 99, column 1, paragraph 2 -----	1-17



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number  
**WO 01/08702 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 39/21**

[HU/US]; 10834 Margate Road, Silver Spring, MD 20901 (US).

(21) International Application Number: **PCT/US00/20641**

(22) International Filing Date: **27 July 2000 (27.07.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
60/146,240 28 July 1999 (28.07.1999) US  
60/178,989 28 January 2000 (28.01.2000) US  
60/200,445 28 April 2000 (28.04.2000) US

(71) Applicant (for all designated States except US): **THE GOVERNMENT OF THE UNITED STATES OF AMERICA**, represented by **THE SECRETARY OF THE DEPARTMENT OF HEALTH AND HUMAN SERVICES** [US/US]; Bethesda, MD 20892 (US).

(74) Agents: **LOCKYER, Jean, M.** et al.; Townsend and Townsend and Crew LLP, 8th Floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FRANCHINI, Genoveffa** [US/US]; 4400 17th Street N.W., Washington, DC 20011 (US). **HEL, Zdenek** [CZ/US]; 11120 Troy Road, North Bethesda, MD 20852 (US). **SHEARER, Gene** [US/US]; 5512 Glenwood Road, Bethesda, MD 20817 (US). **TARTAGLIA, James** [US/US]; 7 Christina Drive East, Schenectady, NY 12303 (US). **NACSA, Janos**

(88) Date of publication of the international search report:  
30 August 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **IMMUNOTHERAPY IN HIV INFECTED PERSONS USING VACCINES AFTER MULTI-DRUG TREATMENT**

(57) Abstract: This invention relates to an improved method of maintaining an immuno-protective response in persons infected with a retrovirus after highly active anti-retroviral therapy.

WO 01/08702 A3

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/20641

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K39/21

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 08539 A (CHIRON CORPORATION) 5 March 1998 (1998-03-05) the whole document ---	1-17
X	M. JOHN ET AL.: "Control of HIV replication by cytotoxic T-lymphocyte responses" J. HIV THERAPY, vol. 4, no. 4, 1999, pages 91-87, XP000982254 page 95, column 1, paragraph 3 -page 96 ---	1-17
X	F. GOTCH ET AL.: "Therapeutic Vaccines in HIV 1 infection" IMMUNOLOGICAL REVIEWS, vol. 170, 1999, pages 173-182, XP000982295 page 179, column 2; figure 1 --- -/--	1-17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

16 March 2001

Date of mailing of the international search report

26/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Skelly, J

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/20641

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>R. MACGREGOR ET AL.: "First human trial of a DNA-based vaccine for treatment of HIV1 infection: safety and host response" J. INF. DIS., vol. 178, 1998, pages 92-100, XP000982247 page 99, column 1, paragraph 2</p>	1-17

# INTERNATIONAL SEARCH REPORT

### Information on patent family members

International Application No.

PCT/US 00/20641

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9808539 A	05-03-1998	AU 4088697 A EP 0942747 A	19-03-1998 22-09-1999

# PATENT COOPERATION TREATY

# PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>15280-4003PC</b>	<b>FOR FURTHER ACTION</b>		see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.
International application No. <b>PCT/US 00/ 20641</b>	International filing date (day/month/year) <b>27/07/2000</b>	(Earliest) Priority Date (day/month/year) <b>28/07/1999</b>	
Applicant <b>THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as</b>			

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 4 sheets.  
☒ It is also accompanied by a copy of each prior art document cited in this report.

**1. Basis of the report**

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.
- ☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).
- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :
- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

**4. With regard to the title,**

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established by this Authority to read as follows:

**5. With regard to the abstract,**

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

**6. The figure of the drawings to be published with the abstract is Figure No.**

- ☐ as suggested by the applicant.
- ☐ because the applicant failed to suggest a figure.
- ☐ because this figure better characterizes the invention.

☒ **None of the figures.**

# INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 00/20641

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K39/21

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)  
BIOSIS, EPO-Internal, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 08539 A (CHIRON CORPORATION) 5 March 1998 (1998-03-05) the whole document ---	1-17
X	M. JOHN ET AL.: "Control of HIV replication by cytotoxic T-lymphocyte responses" J. HIV THERAPY, vol. 4, no. 4, 1999, pages 91-87, XP000982254 page 95, column 1, paragraph 3 -page 96 ---	1-17
X	F. GOTCH ET AL.: "Therapeutic Vaccines in HIV 1 infection" IMMUNOLOGICAL REVIEWS, vol. 170, 1999, pages 173-182, XP000982295 page 179, column 2; figure 1 --- -/--	1-17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

16 March 2001

Date of mailing of the international search report

26/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Skelly, J

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/20641

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>R. MACGREGOR ET AL.: "First human trial of a DNA-based vaccine for treatment of HIV1 infection: safety and host response" J. INF. DIS., vol. 178, 1998, pages 92-100, XP000982247 page 99, column 1, paragraph 2</p> <p>-----</p>	1-17

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/20641

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

WO 9808539	A	05-03-1998	AU 4088697 A	19-03-1998
			EP 0942747 A	22-09-1999

---